CLENPIQ Instructions for Colonoscopy

2 days before the procedure drink 1 bottle of Magnesium Citrate at 7:00 PM.

- The day before the procedure, you may have only clear liquids. NO SOLID FOODS (See enclosed Clear Liquid Diet Instructions). You will take bowel prep in the evening.
- First dose of the prep is at 6:00 PM. Drink one bottle of Clenpiq followed by 5 8-ounce cups of clear liquids. Continue clear liquid diet.
- Take the second dose of the prep at 10:00 PM. Drink the other bottle of Clenpiq followed by 4 8-ounce cups of clear liquids. Continue clear liquid diet until midnight. Then nothing by mouth until after your procedure.
- If you experience vomiting, stop drinking the prep for 20-30 minutes then restart and finish the remainder of the prep. If you are unable to finish the remainder of the bowel prep due to vomiting please call to office.
- On the day of the procedure, do not eat/drink anything before the procedure.

Please take any blood pressure, thyroid or seizure medications with just enough water to swallow them.

*Stop all supplements including fish oil, iron, multi-vitamins, etc. 4 days before the procedure. *

Do not take diabetic medication the morning of your procedure

Appointment Information

Please arrive 30 minutes early

Date:	
Time:	

If you cannot keep your appointment, please call to cancel this one and you may arrange another appointment at that time. Questions? (210) 509-8888

Procedure Locations:

- Theda Oaks Surgery Center #103: 19226 Stone Hue, San Antonio TX 78258 (210-268-0100)
- Stone Oak Methodist Hospital (admitting): 1139 E. Sonterra Blvd, San Antonio TX 78258 (210-638-2000)
- North Central Baptist Hospital (admitting): 520 Madison Oak, San Antonio TX 78258 (210-297-4000)
- Christus Santa Rosa Hospital Westover Hills: 11212 State Hwy. 151 San Antonio TX 78251

How to Prepare for Your Colonoscopy

On the day before your procedure...

- You will be on clear liquids ONLY; NO SOLID FOOD should be consumed the day before
 or the morning of your colonoscopy.
 - DO NOT eat or drink anything that is RED or PURPLE or any DAIRY PRODUCTS

Any of the following items are ok to consume/have.



Water



Gatorade, Powerade, Pedialyte, Sodas, non-carbonated soft drinks like kool-aid. (No red or purple drinks)



Strained fruit juices (without pulp), lemonade, limeade, apple juice, orange juice, white grape juice, or white cranberry juice



Coffee and tea with or without sweetener/sugar. **NO Creamer or Half & Half**



clear broth or bouillon



Jello or Gelatin without fruit or toppings (NOTHING Red or Purple)



Popsicles (No red or purple and No Ice cream center)

Medications to Avoid Prior to Procedure

Please STOP the following medications 2 to 3 days prior to your procedure date:

	Λ Λ	lvil
_	$ \sim$ \sim	ıvıı

Aleve

Alka Seltzer

Anaprox

Ansaid

Arthrotec

Aspirin

Brilinta

Chondroitin

Clinoril

Daypro

Diclofenac

Ecotrin

Eliquis

Etodolac

Feldene

Ginko Biloba

Glucosamine

Ibuprofen

Indomethacin

Ketorolac

• Lodine

Meloxicam

Mobic

Motrin

Nabumetone

Naprosyn

Naproxen

Peroxicam

• Pradax

Relafen

Salsalate

Sulindac

Toradol

• Trental

Trillsate

Voltaren

Xarelto

Please **STOP** the following medications **5 days prior** to your procedure date:

Warfarin

Please STOP the following medications 7 days prior to your procedure date:

- Plavix
- Effient

THEDA OAKS ENDOSCOPY CENTER GUIDELINES 210-268-0100 ext 0

• Things to Remember:

- Opt in to CaseTabs to receive information from Surgery Center
- o For Colonoscopies:
 - Clear Liquids diet the day before the procedure.
 - The more fluids you drink the cleaner you will be for the procedure.
 - Notify your provider if bowel movements are not clear after bowel prep

• Things to Avoid:

- No Recreational drugs, including marijuana, for at least 24 hours prior to procedure
- For Colonoscopies
 - Nothing at all to drink at least 2 hours prior to arrival time (includes gum, hard candy, or sips of water)
- For Endoscopies:
 - Nothing to eat or drink after midnight the day of the procedure (includes gum, hard candy, or sips of water)

Medications:

- Ok to take Blood Pressure, antiarrhythmic, thyroid, or seizure medication on day of procedure with sip of water at least 2 hours prior to arrival time.
- o Please do not take any diabetic medication the day of the procedure.
- o If you are taking any blood thinning medication:
 - Please contact provider's office for further instructions.
 - Contact office to see Cardiac Clearance is needed.

Items to bring with you:

- O Please bring a list of medications including dosage and frequency
 - i.e Lisinopril 20mg daily
- Insurance Card
- Picture ID / ID Card
- Method of Payment (in case of co-pay)
- Warm socks
- You will need a driver Patients who received anesthesia are only allowed to leave the facility in the company of a responsible adult.

Covid Precautions:

- When to reschedule:
 - If you or anyone you have had contact with has had Covid within 14 days
 - If you or anyone you have had contact with is pending the results of a Covid test
 - If you or anyone you have had contact with has had any symptoms of Covid
 - i.e. Fever (100* F), Cough, Muscle aches, Loss of taste / smell ect.
- Day of Procedure:
 - We ask that you please wear a mask to the facility
 - Your temperature and Oxygen Saturation will be taken by Front Desk Staff
 - Only the patient will be allowed to wait in the waiting room
 - We ask that the driver either stay in the car or provide a number to be reached at when the patient arrives in Recovery